



Metabolic responses to temperature stress under elevated pCO₂ in *Crepidula fornicata*

Fanny Noisette, Joelle Richard, Ines Le Fur, Lloyd S. Peck, Dominique Davoult, Sophie Martin

► To cite this version:

Fanny Noisette, Joelle Richard, Ines Le Fur, Lloyd S. Peck, Dominique Davoult, et al.. Metabolic responses to temperature stress under elevated pCO₂ in *Crepidula fornicata*. *Journal of Molluscan Studies*, 2015, 81 (2), pp.238-246. 10.1093/mollus/eyu084 . hal-01100959

HAL Id: hal-01100959

<https://hal.sorbonne-universite.fr/hal-01100959>

Submitted on 9 Jan 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Title: METABOLIC RESPONSES TO TEMPERATURE STRESS UNDER
ELEVATED $p\text{CO}_2$ IN THE SLIPPER LIMPET *CREPIDULA FORNICATA***

NOISETTE F^{*}, RICHARD J, LE FUR I, PECK LS, DAVOULT D, MARTIN S

NOISETTE Fanny (fanny.noisette@sb-roscoff.fr)

LE FUR Ines (Ines.LEFUR@eaurmc.fr)

DAVOULT Dominique (davoult@sb-roscoff.fr)

MARTIN Sophie (sophie.martin@sb-roscoff.fr)

*1 Sorbonne universités, UPMC Univ Paris 06, UMR 7144, Station Biologique de Roscoff,
Place Georges Teissier, 29680 Roscoff Cedex, France*

*2 CNRS, UMR 7144, Station Biologique de Roscoff, Place Georges Teissier, 29680 Roscoff
Cedex, France*

RICHARD Joëlle (Joelle.Richard@univ-brest.fr)

*3 Université de Bretagne Occidentale, Institut Universitaire Européen de la Mer, Laboratoire
des Sciences de l'Environnement Marin (UMR CNRS 6539), Technopôle Brest-Iroise, Place
Copernic, F-29280 Plouzané, France.*

*4 Natural Environment Research Council British Antarctic Survey, High Cross, Madingley
Road, Cambridge CB3 0ET, United Kingdom*

PECK Lloyd S. (lspe@bas.ac.uk)

*4 Natural Environment Research Council British Antarctic Survey, High Cross, Madingley
Road, Cambridge CB3 0ET, United Kingdom*

Short running head: *C. fornicata* respiration under high $p\text{CO}_2$

* Corresponding author: Fanny NOISETTE

Email: fanny.noisette@sb-roscoff.fr

Postal address: Station Biologique de Roscoff, Place Georges Teissier, 29 680 ROSCOFF (France)

Phone number: +33 298292333

ABSTRACT

In the current context of environmental change, ocean acidification is predicted to affect the cellular processes, physiology and behavior of all marine organisms, impacting survival, growth and reproduction. In relation to thermal tolerance limits, the effects of elevated $p\text{CO}_2$ could be expected to be more pronounced at the upper limits of the thermal tolerance window. Our study focused on *Crepidula fornicata*, an invasive gastropod which colonized shallow waters around European coasts during the 20th century. We investigated the effects of 10 weeks' exposure to current (380 μatm) and elevated (550, 750, 1000 μatm) $p\text{CO}_2$ on this engineer species using an acute temperature increase ($1^\circ\text{C } 12\text{h}^{-1}$) as the test. Respiration rates were measured on both males (small individuals) and females (large individuals). Mortality increased suddenly from 34°C , particularly in females. Respiration rate in *C. fornicata* increased linearly with temperature between 18°C and 34°C , but no differences were detected between the different $p\text{CO}_2$ conditions either in the regressions between respiration rate and temperature, or in Q_{10} values. In the same way, condition indices were similar in all the $p\text{CO}_2$ treatments at the end of the experiment but decreased from the beginning of the experiment. This species was highly resistant to acute exposure to high temperature regardless of $p\text{CO}_2$ levels, even though food was limited during the experiment. *C. fornicata* appears to have either developed resistance mechanisms or a strong phenotypic plasticity to deal with fluctuations of physico-chemical parameters in their habitat. This suggests that this invasive species may be more resistant to future environmental changes compared to its native competitors.

Keywords: CO_2 stress, invasive species, ocean acidification, Q_{10} , respiration, temperate waters

INTRODUCTION

As part of global change, ocean acidification is caused by increasing anthropogenic CO₂ emissions which have increased since the beginning of the industrial revolution (Solomon *et al.*, 2007). Future *p*CO₂ increases are predicted to reduce the pH of surface waters by 0.3 - 0.4 units by the end of the century (Caldeira & Wickett, 2003). Such decreases will produce changes in carbon and carbonate seawater chemistry through decreased carbonate ion concentrations (CO₃²⁻) and a lower calcium carbonate saturation state (Ω). These changes are predicted to have major consequences for marine life (Fabry *et al.*, 2008; Kroeker *et al.*, 2013b) and, especially, could have broad impacts on physiological functions of heterotrophic marine organisms (Pörtner, 2008; Hofmann & Todgham, 2010).

The decrease in pH is likely to have a wide range of effects on marine invertebrates via shifts in acid-base homeostasis, changes in metabolism and energy balance (Pörtner *et al.*, 2005), leading to effects on somatic growth (Berge *et al.*, 2006; Thomsen & Melzner, 2010), respiration (Melatunan *et al.*, 2011; Schalkhausser *et al.*, 2013), excretion (Liu & He, 2012), calcification (Gazeau *et al.*, 2007; Wood *et al.*, 2008; Watson *et al.*, 2012) or feeding rates (Bamber, 1990; Navarro *et al.*, 2013). Many marine invertebrates exposed to elevated *p*CO₂ have exhibited metabolic depression (Willson & Burnett, 2000; Michaelidis *et al.*, 2005; Navarro *et al.*, 2013) as a decrease in respiration rate while others have remained unaffected (Gutowska *et al.*, 2008; Lannig *et al.*, 2010; Clark *et al.*, 2013) or even increased their metabolic rate (Wood *et al.*, 2008; Beniash *et al.*, 2010). These responses are highly species-specific and may vary with organism size (Beniash *et al.*, 2010). The resilience of the species studied, and the capacity to regulate metabolism under stressful conditions are also important (Pörtner, 2008). These physiological impacts are likely to have broad effects on the survival, growth and reproduction of marine species (Shirayama & Thornton, 2005; Byrne, 2011),

which would lead to changes in community structure from altered diversity and abundances (Hale *et al.*, 2011; Kroeker *et al.*, 2013a).

These physiological impacts are likely modulated by temperature because temperature is a primary driver of physiological function in ectotherms (Hofmann & Todgham, 2010). Increasing temperature affects the rate of all biochemical reactions, and hence cellular processes and physiological functions (Clarke, 1983; Pörtner, 2012), increasing metabolic costs within a limited thermal tolerance window (Peck *et al.*, 2002; Marshall *et al.*, 2003). The interactive effects of increased temperature and elevated CO₂ concentrations are predicted to impair physiological processes (Clarke, 2003; Pörtner, 2008) by narrowing the thermal tolerance window of the organisms (Metzger *et al.*, 2007; Lannig *et al.*, 2010) and elevating vulnerability to extreme temperature (Schalkhauser *et al.*, 2012).

In a context of global change, non-indigenous species are expected to be favored in their introduced area (Dukes & Mooney, 1999; Occhipinti-Ambrogi, 2007) mainly because robustness to abiotic variation is often a trait that determines the success of invasive of a species (Hellmann *et al.*, 2008; Lenz *et al.*, 2011). Climatic changes in the physical environment will likely affect the distribution, spread, abundance, impacts and interactions of species, possibly to the advantage of introduced organisms (Occhipinti-Ambrogi, 2007). Thus our study focused on the response of an invasive Calyptraeidae gastropod living on western European coasts, but which originates from North East America. The slipper limpet, *Crepidula fornicata* (Linné 1758) was introduced in Europe at the end of the 19th century, mainly with oysters (*Crassostrea gigas*) which were imported for farming (Blanchard, 1995), and has subsequently colonized European coasts from southern Sweden to southern France (Blanchard, 1997). *C. fornicata* has significant impacts on biodiversity and ecosystem functioning where it has established (De Montaudouin *et al.*, 1999; Decottignies *et al.*, 2007; Martin *et al.*, 2007). It lives in shallow sites, especially in bays and estuaries where very high

densities of over one thousand individuals m^{-2} have been reported (Blanchard, 1995). *C. fornicata* is known to be strongly resistant to environmental variations, particularly temperature and salinity (Blanchard, 1995; Blanchard, 1997; Diederich & Pechenick, 2013). In light of the different ecological and physiological characteristics of *C. fornicata*, it is important to investigate the impact of future $p\text{CO}_2$ levels, and determine its resistance capacities to high levels of stress to assess the likely future impact of this engineer species in the ecosystems to which it was introduced.

The present study was designed to investigate the metabolic responses of *C. fornicata* to high $p\text{CO}_2$ conditions during temperature stress. Short-term experimental approaches using faster temperature elevations than natural changes provide valuable insight into physiological responses of marine invertebrates in term of their ability to resist high levels of stress or their lethal temperature (Sokolova & Pörtner, 2003; Peck *et al.*, 2004; Pörtner *et al.*, 2006; Richard *et al.*, 2012). Following the hypothesis that CO_2 stress will increase sensitivity to temperature change, we evaluated changes in oxygen-consumption of *C. fornicata* individuals previously reared under elevated $p\text{CO}_2$ for 10 weeks during a rapid temperature increase ($1^\circ\text{C } 12\text{h}^{-1}$). Respiration rates were measured as a proxy for metabolism on males (small individuals) and females (large individuals), as in this species there is sexual dimorphism in size.

MATERIAL & METHODS

Biological material

Crepidula fornicata stacks were collected by SCUBA divers on 4 February 2010, in Morlaix Bay (northwest Brittany, France), at the “Barre des Flots” site ($3^\circ 53.015'\text{W}$; $48^\circ 40.015'\text{N}$) at a depth of 10 meters and at an *in situ* temperature of 11.6°C (SOMLIT: *Service d’Observation de la Mer et du Littoral* data). They were transferred directly to

aquaria at the Station Biologique de Roscoff where they were held in natural unfiltered seawater at a temperature around 10°C, until they were used in experiments starting on 10 March 2010.

Males and females at the top and the bottom of stacks respectively, were selected, separated and individually labelled. Small males (23.31 ± 0.16 mm length), which were still slightly mobile, were placed individually on 3 cm Petri dishes one month before the beginning of the trials. Dead individual shells at the base of stacks were kept as the substratum under the largest living immobile females (47.53 ± 0.25 mm length). In *C. fornicata*, size cannot be discriminated from sex because this is a protandrous hermaphroditic organism, changing sex with age and size (Coe 1938). All individuals were gently brushed to remove epibionts and biofilm from their shells before proceeding to the metabolic measurements.

Condition indices (CI) were calculated on a pool of 20 specimens in March, before the beginning of the experiment, and on all remaining living and recently dead individuals (male $n = 74$; female $n = 99$) at the end of the temperature increase on 29 May 2010. Shell dry weight (DW_{Shell}), shell length and tissue dry weight (DW_{Tissue}) were determined separately on each individual after drying at 60°C for 48h. Specimens were then ignited in a muffle furnace at 520°C for 6 h, with tissue ash-free dry weight ($AFDW_{\text{Tissue}}$) being obtained by difference. CI were calculated as:

$$CI = (AFDW_{\text{Tissue}} / DW_{\text{Shell}}) \times 100.$$

Mortality was checked daily during the experiment. Individuals with no reaction when the foot was stimulated were classed as dead and removed from the tanks.

Experimental conditions and set-up

After distributing randomly in each of twelve 10-L aquarium tanks comprising the experimental flow-through system (as described in Noisette *et al.*, 2013), 120 males and 120

females (i.e. 10 individuals of each sex per aquarium) were held in different $p\text{CO}_2$ conditions between 13 March and 29 May 2010. At the beginning of the experiment, pH was gradually decreased (by bubbling CO_2) over four days at $0.1 \text{ pH units day}^{-1}$ from 8.1 until the required pH was reached. Specimens were subsequently held for ten weeks in four different $p\text{CO}_2$ conditions: a current $p\text{CO}_2$ of $380 \text{ } \mu\text{atm}$ ($\text{pH}_\text{T} = 8.07$), and three elevated $p\text{CO}_2$ levels of $550 \text{ } \mu\text{atm}$ ($\text{pH}_\text{T} = 7.94$), $750 \text{ } \mu\text{atm}$ ($\text{pH}_\text{T} = 7.82$) and $1000 \text{ } \mu\text{atm}$ ($\text{pH}_\text{T} = 7.77$). The elevated $p\text{CO}_2$ values corresponded to different scenarios predicted by the Intergovernmental Panel on Climate Change (IPCC) for the end of the century (Solomon *et al.*, 2007) and were selected according to the recommendations of Barry *et al.*, (2010). $p\text{CO}_2$ was adjusted by bubbling CO_2 -free air (current $p\text{CO}_2$) or pure CO_2 (elevated $p\text{CO}_2$) in four 100 L header tanks (1 per $p\text{CO}_2$ condition) supplied with natural unfiltered seawater pumped from the sea, directly at the foot of the Station Biologique de Roscoff. Seawater was continually delivered by gravity from each header tank to three aquaria per $p\text{CO}_2$ condition at a constant rate of 9 L h^{-1} (renewal rate: 90% total aquarium volume h^{-1}). $p\text{CO}_2$ was monitored and controlled by a feedback system (IKS Aquastar, Karlsbad, Germany) that regulated the addition of gas in the header tanks. pH values of the pH-stat system were adjusted from daily measurements of pH on the total scale (pH_T) in the aquaria using a pH meter (HQ40D, Hach Lange, Ltd portable LDOTM, Loveland, Colorado, USA) calibrated using Tris/HCl and 2-aminopyridine/HCl buffers (Dickson *et al.*, 2007). The twelve aquaria were placed in four thermostatic baths where temperature was controlled to $\pm 0.2 \text{ }^\circ\text{C}$ using 100 - 150 W submersible heaters.

Before the rapid temperature increase experiment, *C. fornicata* individuals were maintained in the different $p\text{CO}_2$ treatments for 10 weeks while temperature was raised successively to mimic the natural rate of temperature change from winter to summer. Temperature was maintained at 10°C from the beginning of the trial to 29 March. It was raised to 13°C from 5 to 19 April and to 16°C from 26 April to 18 May 2010. To reach these

set levels the temperature was increased by $0.5^{\circ}\text{C day}^{-1}$ until the new set temperature was achieved. During the experiment, animals were naturally fed by the phytoplankton provided by unfiltered seawater.

The rapid temperature increase experiment was conducted between the 18 and 29 May 2010. In all four $p\text{CO}_2$ treatments, temperature was increased from 16 to 36°C at $1^{\circ}\text{C } 12\text{h}^{-1}$. *C. fornicata* oxygen consumption was measured (see below) both in small and large individuals in the different $p\text{CO}_2$ treatments during this rapid temperature increase.

Seawater parameters

Seawater parameters were monitored throughout the experiment. pH_T and temperature were recorded daily in each of the 12 aquaria using a pH meter (HQ40D, Hach Lange, Ltd portable LDOTM, Loveland, Colorado, USA). Total alkalinity was determined every 3 weeks by 0.01N HCl potentiometric titration on an automatic titrator (Titroline alpha, Schott SI Analytics, Mainz, Germany). Seawater carbonate chemistry, *i.e.* exact CO_2 partial pressure ($p\text{CO}_2$) and saturation state of aragonite were calculated in each $p\text{CO}_2$ condition using CO_2SYS software (Lewis & Wallace, 1998) using constants from Mehrbach *et al.*, (1973) refitted by Dickson & Millero, (1987). Mean values (\pm standard error, SE) of the parameters in each $p\text{CO}_2$ treatment are presented in Table 1.

Oxygen consumption measurements

During the rapid temperature increase trial (18 - 29 May 2010), oxygen consumption of 6 randomly selected labeled individuals of each sex (2 per aquaria) was measured in each of the $p\text{CO}_2$ treatments every two days, at 18, 22, 26, 30 and 34°C . Respiration rates were determined using closed incubations in 75 mL (males) or 180 mL (females) acrylic chambers (Engineering & Design Plastics Ltd, Cambridge, UK) filled with water from the same

aquarium (see methods in Morley *et al.*, 2007). Chambers were placed in their respective aquaria during incubations to keep the temperature constant. Incubations varied between 1 h and 3 h depending on temperature and were halted before oxygen saturation fell below 80% saturation. Control incubations without animals (n = 1 control incubation / aquarium / measurement) were carried out to allow correction for microbial activity in seawater.

Respiration rates were calculated from the differences in measurements of oxygen concentration during trials and controls using a non-invasive fiber-optical system (FIBOX 3, PreSens, Regensburg, Germany) made up of an optical fiber and reactive oxygen spots attached to the inner wall of the chambers. These spots were calibrated with 0% and 100% oxygen buffers made from the manufacturer instructions. 0% O₂ buffer was prepared by dissolving 10 g of Na₂SO₃ in 1 L of seawater and 100% O₂ buffer was prepared by bubbling air in 1L of seawater for 20 min to achieve oxygen saturation. Previous experiments had demonstrated that oxygen consumption remained linear during all the incubation periods. Chamber contents were mixed gently by inverting chambers several times before each oxygen measurement. Respiration (R) rates (in $\mu\text{mol O}_2 \text{ g}^{-1} \text{ AFDW h}^{-1}$) were corrected for oxygen consumption in controls and calculated as:

$$R = -(\Delta\text{O}_2 \times V) / (\Delta t \times \text{AFDW}_{\text{Tissue}})$$

where ΔO_2 ($\mu\text{mol O}_2 \text{ L}^{-1}$) is the difference between initial and final O₂ concentrations during the incubation, V (L) is the chamber volume minus the individual *C. fornicata* volume, Δt (h) is the incubation time and $\text{AFDW}_{\text{Tissue}}$ (g) is the tissue ash free dry weight of the slipper limpet incubated.

Q₁₀ coefficients were calculated by using the standard equation:

$$Q_{10} = (R_H / R_L)^{10 / (T_H - T_L)}$$

where T_L and T_H were the lowest and highest temperature reached and R_L and R_H the respiration rates in these temperature respectively.

Statistical analyses

All statistical analyses were performed using R version 2.15.0 (R Core Team 2013) and STATISTICA software. A logistic regression (general linear model, GLM) was applied to test the differences in mortalities between the different $p\text{CO}_2$ treatments and between sex with temperature as the linear variable. The effects of $p\text{CO}_2$, sex and the interaction of these two factors on condition index (CI) at the end of the experiment and on Q_{10} values were investigated by 2-way analysis of variance (ANOVA). Linear regressions between respiration rates and increasing temperatures were fitted in the four different $p\text{CO}_2$ treatments for males and females separately. Differences between $p\text{CO}_2$ treatments were explored using an ANCOVA with $p\text{CO}_2$ and sex as fixed factors and temperature as co-variable.. Normality was assessed using the Kolmogorov-Smirnov test and Levene's test was used to ensure that variances were homogenous. All the results are presented as mean \pm standard error (SE).

RESULTS

Mortality occurred between 34 and 36°C for females and 22 and 36°C for males (Figure 1). There were no significant differences in mortality between the different $p\text{CO}_2$ treatments (GLM, $df = 3$, $F = 0.680$, $p = 0.565$) or between males and females (GLM, $df = 1$, $F = 0.580$, $p = 0.449$). Moreover, the interaction between factors $p\text{CO}_2$ and sex of the individuals was not significant (GLM, $df = 3$; $F = 0.21$; $p = 0.888$). At $p\text{CO}_2$ levels of 380, 550, 750 and 1000 μatm , the mortality was 29, 19, 19, and 24 for females and 28, 6, 8, and 6 for males . At the end of the acute temperature increase nearly twice the number of females had died (91) compared with the males (48) (χ^2 test, $p < 0.05$).

The mean condition index before the start of the experiment was 3.00 ± 0.27 ($n=10$). It varied at the end of the experiment between 1.69 ± 0.13 for males at $p\text{CO}_2$ of 380 μatm and 2.41 ± 0.27 for females at $p\text{CO}_2$ of 550 μatm (Table 2). There were no effects of $p\text{CO}_2$, sex or the interaction of these two factors on the condition index at the end of the trial (Table 2). However, the condition index from the beginning of the experiment (3.00 ± 0.27) was different from the mean condition index including all $p\text{CO}_2$ conditions (2.11 ± 0.07) at the end of the trial (t-test, $df = 181$, $t = 3.159$, $p = 0.002$), which means that CI in both males and females decreased significantly from the start to the end of the experiment (Figure 2).

Female respiration rates varied between $0.51 \mu\text{mol O}_2 \text{ g}^{-1} \text{ AFDW h}^{-1}$ at 18°C and $p\text{CO}_2$ of 750 μatm and $91.62 \mu\text{mol O}_2 \text{ g}^{-1} \text{ AFDW h}^{-1}$ at 32°C and $p\text{CO}_2$ of 380 μatm . Males had higher rates, which ranged between $5.13 \mu\text{mol O}_2 \text{ g}^{-1} \text{ AFDW h}^{-1}$ at 18°C and $p\text{CO}_2$ of 380 μatm and $175.51 \mu\text{mol O}_2 \text{ g}^{-1} \text{ AFDW h}^{-1}$ at 32°C and $p\text{CO}_2$ of 380 (Figure 3).

Relationships between respiration rate and temperature were linear at each $p\text{CO}_2$ level (Figure 3). Respiration rose significantly with increasing temperature in all $p\text{CO}_2$ treatments, for both males and females (Table 3, all p -values < 0.02). There were no significant differences between the slopes of the different regressions among the $p\text{CO}_2$ treatments or between sexes (analysis of slopes, $df = 3$, $F = 1.1$, $p = 0.346$). The intercepts of the different regressions also did not significantly vary among $p\text{CO}_2$ (ANCOVA, $df = 3$, $F = 0.350$, $p = 0.789$), but there were difference between males and females (ANCOVA, $df = 1$, $F = 62.63$, $p < 0.001$).

Q_{10} values ranged from 1.24 to 2.40 for females and from 1.36 to 2.77 for males among the different $p\text{CO}_2$ treatments (Figure 2). There was no significant $p\text{CO}_2$ effect on Q_{10} values for either males or females (Table 2). Across all $p\text{CO}_2$ treatments, females had significantly lower Q_{10} values than males with means of 1.61 ± 0.11 and 2.00 ± 0.12 for

females and males, respectively (Table 2). The interaction between $p\text{CO}_2$ and sex, however, was not significant (Table 2).

DISCUSSION

Independently of the impact of $p\text{CO}_2$ we planned to test, one of the major issues of this study was food limitation which was unintentionally imposed on the *C. fornicata* individuals in the experiments. This food limitation was detected because the decrease in condition indices (CI) of both males and females from the beginning to the end of the experiment. Such decreases in CI are usually related to food quantity or quality supplied to organisms (Norkko & Thrush, 2006). Animals were maintained in unfiltered seawater which carried natural phytoplankton at a concentration between 0.2 and 1 $\mu\text{g Chl a L}^{-1}$ (SOMLIT data). The water renewal in the aquarium was maintained constant at a rate of 0.9 L h⁻¹ (i.e. 90% of the total volume of each aquarium changed per hour). Water supply in our experimental system was likely too low to provide sufficient food for the experimental animals, which thus relied on internal energy reserves and so decreased their CI. A similar outcome was reported for mussels by Mackenzie *et al.* (2014).

The use of stored reserves was similar in the different $p\text{CO}_2$ conditions as CI at the end of the experiment did not differ between the different $p\text{CO}_2$ treatments, and this was the case for both sexes. Previous studies have shown interspecific variability in the responses of condition indices under high $p\text{CO}_2$ levels, ranging from a lack of effect (Cummings *et al.*, 2011; Clark *et al.*, 2013; Sanders *et al.*, 2013) to large changes in condition under high $p\text{CO}_2$ levels (Hiebenthal *et al.*, 2013; Range *et al.*, 2014). Energy availability is a major component in mitigating the effects of ocean acidification (Pansch *et al.*, 2014). Studies have shown that an abundant food supply might counteract even overcome the negative effects of high $p\text{CO}_2$

on adult and juvenile bivalves (Melzner *et al.*, 2011; Thomsen *et al.*, 2013). Thus, it is important to consider that in this study *C. fornicata* were under limited food conditions when interpreting their metabolic responses to elevated $p\text{CO}_2$ conditions during the temperature rise. The data here are representative of conditions where there is temperature stress and food supplies are limited, conditions that can occur in the field.

The limitation of food supply was not markedly more important in any of our reduced pH conditions as there were no differences in mortality rates between the different $p\text{CO}_2$ treatments in *C. fornicata* males and females. This is a different outcome to that reported for some other mollusk species held in elevated $p\text{CO}_2$ levels (Shirayama & Thornton, 2005; Beniash *et al.*, 2010). However, similarly to our study, Pansh *et al.*, (2014) showed that food availability had no impact on mortalities of the barnacle *Amphibalanus improvises* held in different $p\text{CO}_2$ conditions. In the present study, important mortalities started to occur from 32°C and they became larger at and above 34°C for both males and females. These values are consistent with the upper lethal temperature recorded for *C. fornicata* by Diederich & Pechenick, (2013) in a laboratory study investigating a population from Rhode Island, USA, in which only 40% of the adults survived after a 3 h exposure to 34°C, and all died after a 3 h exposure to 36°C. Mortality was higher in females (larger individuals) than in males (small individuals) even if, male started to die at lower temperatures than females. Similarly, Peck *et al.*, (2009) demonstrated for 14 species that smaller species survived to higher temperatures than large ones when temperature was raised at 1°C day⁻¹, and Peck *et al.*, (2013) showed that juveniles had higher upper temperature limits than adults in 4 species of marine invertebrates at warming rates of 1°C day⁻¹ and 1°C 3days⁻¹. The mechanisms setting temperature limits at acute rates of warming may not be energy availability (Peck *et al.*, 2014) and females, which had more energetic reserves than males, may thus have not had an advantage.

Despite the decreases in CI, mean respiration rates of *C. fornicata* at 18°C and $p\text{CO}_2$ of 380 μatm were 31 and 26 $\mu\text{mol O}_2 \text{ g}^{-1} \text{ AFDW h}^{-1}$ for males and females, respectively, which are close to the middle of the range of *in situ* values reported for wild individuals from the Bay of Brest (Brittany, France) (6 to 63 $\mu\text{mol O}_2 \text{ g}^{-1} \text{ AFDW h}^{-1}$: Martin *et al.*, 2006). This indicates that animals in the experiments here had similar oxygen consumption than wild specimens and were not metabolically depressed under insufficient food supply. In both *C. fornicata* males and females, respiration rates increased with temperature, as previously demonstrated for this species by Newell & Kofoed, (1977) and most ectotherm metabolic rates are correlated positively with temperature (Cossins & Bowler, 1987). Respiration rates were higher in *C. fornicata* males than in females regardless of the temperature. Generally, mass-specific respiration rates of small individuals are higher than those of larger ones because metabolic rate (normalized to the biomass) decreases with increasing organisms size (von Bertalanffy, 1951; Parsons *et al.*, 1984).

The relationship between oxygen consumption and temperature here for *C. fornicata* was similar in all the different $p\text{CO}_2$ treatments. The slopes and intercepts of the regressions were not significantly different across the four $p\text{CO}_2$ conditions which means temperature effect on respiration rate was not affected by the different $p\text{CO}_2$ levels in males or females. In contrast to our results, Lannig *et al.*, (2010) found that an acute temperature rise (1.25°C/12h) caused a more rapid increase in metabolic rate in *Crassostrea gigas* under elevated $p\text{CO}_2$ conditions, and there was a synergistic effect of temperature and $p\text{CO}_2$. The lack of difference in respiration between animals held in different $p\text{CO}_2$ conditions may be related to a stronger ability to up-regulate their metabolism under a temperature stress irrespective of $p\text{CO}_2$. Thus, under warming conditions, *C. fornicata* can generate sufficient energy to cope with any effects of decreased pH (Wood *et al.*, 2010). Q_{10} values were also similar across $p\text{CO}_2$ treatments in both males and females and they were within the expected

range of values recorded for marine invertebrates (Branch *et al.*, 1988; Marshall *et al.*, 2003). Even if *C. fornicata* individuals were food limited, their oxygen consumption remained unaffected by elevated $p\text{CO}_2$. A similar lack of $p\text{CO}_2$ effect was reported for growth and shell strength of the barnacle *A. improvisus* (Pansch *et al.*, 2014). In our study, the low food supply did not appear to affect the resistance or resilience of *C. fornicata* to CO_2 stress.

Several studies investigating the response of mollusk respiration to elevated $p\text{CO}_2$ have demonstrated metabolic depression under high $p\text{CO}_2$ in both bivalves and gastropods (Michaelidis *et al.*, 2005; Bibby *et al.*, 2007; Fernandez-Reiriz *et al.*, 2011; Melatunan *et al.*, 2011; Liu & He, 2012; Navarro *et al.*, 2013). Conversely, others observed no $p\text{CO}_2$ effect on mollusk respiration and general metabolism (Gazeau *et al.*, 2007; Marchant *et al.*, 2010; Fernandez-Reiriz *et al.*, 2012; Clark *et al.*, 2013) as reported in our study. In some rare cases, O_2 consumption was reported to increase under high $p\text{CO}_2$ conditions (Wood *et al.*, 2010; Cummings *et al.*, 2011). The effects of high CO_2 concentrations on metabolism appear species-specific and depend on resistance capacities of the organisms (Melzner *et al.*, 2009). It has been widely reported that exposure to environmental high $p\text{CO}_2$ levels leads to changes in homeostasis and extracellular acid-base balance counterbalanced by metabolic depression in many cases (Pörtner *et al.*, 2005; Pörtner, 2008), although it should be noted, as above, that metabolic depression is often not seen in high $p\text{CO}_2$ conditions. Differences in acid-base regulatory capacities by increasing HCO_3^- internal concentrations (Michaelidis *et al.*, 2005; Gutowska *et al.*, 2010) or H^+ excretion (Pörtner *et al.*, 2005) are taxon specific and are more or less effective in mitigating the effects of hypercapnia. It has also been suggested that organisms could maintain low metabolic rates without controlling internal pH by not using pH-sensitive oxygen-binding pigments (Thomsen *et al.*, 2010; Hiebenthal *et al.*, 2013). Such mechanisms may be crucial factors in explaining the observed variation in sensitivities and resistances of marine invertebrates to elevated $p\text{CO}_2$ conditions (Gutowska *et al.*, 2010).

It is important to note here that many of the studies to date on the effects of elevated $p\text{CO}_2$ on organisms are short-term and acute (e.g. Tomanek *et al.*, 2011), not reflecting the long-term trade off in energy balance and physiological changes associated with acclimation of new environmental conditions (Clark *et al.*, 2013). For example, metabolic depression acts as a time-limited compensation strategy to survive unfavorable condition such as high CO_2 concentrations (Guppy & Withers, 1999; Willson & Burnett, 2000). Because *C. fornicata* were held for 10 weeks in the different $p\text{CO}_2$ treatments in this investigation, it is likely there was enough time for them to acclimate to the new pH, and no difference in oxygen consumption was detected between the different $p\text{CO}_2$ conditions. However, the energetic cost likely produced by the negative effects of elevated $p\text{CO}_2$ may either be relatively small, or difficult to maintain over longer time periods. This could be seen in impacts on other physiological processes than respiration (Catarino *et al.*, 2012). For example, Bibby *et al.*, (2008) demonstrated that exposure to hypercapnic conditions may compromise the ability to express an immune response in mussels. They showed that *Mytilus edulis* phagocytosis declined as function of decreased pH. In the same way, Matozzo *et al.*, (2012) showed that elevated $p\text{CO}_2$ and temperature may strongly affect haemocyte functionality in the bivalves *Chamelea gallina* and *Mytilus galloprovincialis*. Other cellular processes have also been shown to be negatively impacted by high CO_2 concentrations, including protein synthesis in the sipunculid *Sipunculus nudus* (Langenbuch *et al.*, 2006) or enzyme activities in *C. gallina* and *M. galloprovincialis* (Matozzo *et al.*, 2013). However, studies of the impact of reduced pH on immune systems have generally been of short duration and it would be interesting to investigate other physiological parameters than respiration (e.g. calcification, protein production, immunity regulation, fertility) in *C. fornicata* acclimated over several months in the different $p\text{CO}_2$ conditions predicted for the end of the century. As a coastal species adapted to relatively large fluctuations of abiotic parameters, *C. fornicata* in this study were

strongly resistant to both elevated $p\text{CO}_2$ and increased temperature. Indeed, resistance to high $p\text{CO}_2$ levels can also come from pre-acclimation or pre-adaptation to fluctuations in the environment where species live (Burnett, 1997). Species living in environments with large abiotic variation have a high phenotypic plasticity which can allow them to survive in stressful conditions (Hofmann & Todgham, 2010). Coastal organisms are more exposed to physico-chemical variations than their open-ocean counterparts that live in more stable thermal and pH environments (Berge *et al.*, 2006; Peck *et al.*, 2006). Species living in shallow waters tolerate not only seasonal and extreme temperature events but also periodic large fluctuations in seawater pH, driven by biological process that sequester and release large amounts of CO_2 (Beniash *et al.*, 2010). This exposure to a wide environmental variation has likely led to the evolution of resistance mechanisms to abiotic factors including variations in $p\text{CO}_2$ and/or pH (Lannig *et al.*, 2010).

C. fornicata is an invasive species which has successfully colonized European coastal shallow waters. This species is likely to have high phenotypic plasticity and resilience to physico-chemical variations that determined its success. Indeed, successful invasive species generally share characteristics that allow them to establish, colonize and expand their range. Among these characteristics, tolerance to environmental stress is one of the most common (Lenz *et al.*, 2011). In a global change context, the movement of physico-chemical conditions away from the optimum increases the energy required by marine species to fuel the extra processes entrained to resist the stresses involved and to maintain homeostasis. This may result in changes in overall physiological condition (Cummings *et al.*, 2011) that could impact ecological processes and community interactions. The high resilience to altered $p\text{CO}_2$ /low pH levels observed here for *C. fornicata* may confer a competitive advantage to this invasive species over taxonomically or functionally related species (Lenz *et al.*, 2011). For example, the performance of the scallop *Pecten maximus*, which is one of the *C. fornicata* competitors

(Thouzeau *et al.*, 2000; Fresard & Boncoeur, 2006), has been shown to be negatively affected by high $p\text{CO}_2$ levels (Schalkhauser *et al.*, 2013). These different sensitivities to environmental factors will likely dictate “winners” and “losers” among marine species that could lead to a restructuring of benthic communities. With other studies, our data suggest this restructuring could favor invasive species as evidence is building that shows they are more resistant to change than their native competitors (Dukes & Mooney, 1999; Occhipinti-Ambrogi, 2007).

ACKNOWLEDGMENTS

The authors thank the Marine Operations and Services Department from the Station Biologique de Roscoff for the underwater sampling and the help for system building. This work was supported by the CALCAO project funded from the Region Bretagne, and by the Interreg IVa France (Channel) – England Marinexus project no. 4073 funded by the FEDER programme. It also contributes to the “European Project on Ocean Acidification” (EPOCA) which received funding from the European Community’s Seventh Framework Programme (FP7/2007-2013) under grant agreement n° 211384

REFERENCES

- Bamber R (1990) The effects of acidic seawater on three species of lamellibranch mollusc. *Journal of Experimental Marine Biology and Ecology* **143**: 181-191
- Barry JP, Tyrrell T, Hansson L, Plattner GK, Gattuso JP (2010) Atmospheric CO₂ targets for ocean acidification perturbation experiments. In: Riebesell U, FVJ, Hansson L. & Gattuso J.-P. (ed) Guide to best practices for ocean acidification research and data reporting, Luxembourg: Publications Office of the European Union, pp 260
- Beniash E, Ivanina A, Lieb NS, Kurochkin I, Sokolova IM (2010) Elevated level of carbon dioxide affects metabolism and shell formation in oysters *Crassostrea virginica*. *Marine Ecology-Progress Series* **419**: 95-108
- Berge JA, Bjerkeng B, Pettersen O, Schaanning MT, Øxnevad S (2006) Effects of increased sea water concentrations of CO₂ on growth of the bivalve *Mytilus edulis* L. *Chemosphere* **62**: 681-687
- Bibby R, Cleall-Harding P, Rundle S, Widdicombe S, Spicer J (2007) Ocean acidification disrupts induced defences in the intertidal gastropod *Littorina littorea*. *Biology Letters* **3**: 699-701
- Bibby R, Widdicombe S, Parry H, Spicer J, Pipe R (2008) Effects of ocean acidification on the immune response of the blue mussel *Mytilus edulis*. *Aquatic Biology* **2**: 67-74
- Blanchard M (1995) Origine et état de la population de *Crepidula fornicata* (Gastropoda Prosobranchia) sur le littoral français. *Halietis* **24**: 75-86
- Blanchard M (1997) Spread of the slipper limpet *Crepidula fornicata* (L. 1758) in Europe. Current state and consequences. *Scientia Marina* **61**: 109-118
- Branch GM, Borchers P, Brown CR, Donnelly D (1988) Temperature and food as factors influencing oxygen consumption of intertidal organisms, particularly limpets. *American Zoologist* **28**: 137-146

480 Burnett LE (1997) The challenges of living in hypoxic and hypercapnic aquatic environments.
 481 *American Zoologist* **37**: 633-640

482 Byrne M (2011) Impact of ocean warming and ocean acidification on marine invertebrate life
 483 history stages: vulnerabilities and potential for persistence in a changing ocean. In:
 484 Gibson R, Atkinson R, Gordon J, Smith I, Hughes D (eds) *Oceanography and Marine*
 485 *Biology: An Annual Review*. Taylor & Francis, pp 1-42

486 Caldeira K, Wickett ME (2003) Anthropogenic carbon and ocean pH. *Nature* **425**: 365-365

487 Catarino AI, Bauwens M, Dubois P (2012) Acid-base balance and metabolic response of the
 488 sea urchin *Paracentrotus lividus* to different seawater pH and temperatures.
 489 *Environmental Science and Pollution Research* **19**: 2344-2353

490 Clark MS, Thorne MAS, Amaral A, Vieira F, Batista FM, Reis J, Power DM (2013)
 491 Identification of molecular and physiological responses to chronic environmental
 492 challenge in an invasive species: the Pacific oyster, *Crassostrea gigas*. *Ecology and*
 493 *Evolution* **3**: 3283-3297

494 Clarke A (1983) Life in cold water: the physiological ecology of polar marine ectotherms.
 495 *Oceanography and Marine Biology* **21**: 341-453

496 Coe WR (1938) Influence of association on the sexual phases of gastropods having protandric
 497 consecutive sexuality I. *The Biological Bulletin* **75**: 274-285

498 Cossins AR, Bowler K (1987) Temperature biology of animals. Chapman and Hall London

499 Cummings V, Hewitt J, Van Rooyen A, Currie K, Beard S, Thrush S, Norkko J, Barr N,
 500 Heath P, Halliday NJ, Sedcole R, Gomez A, McGraw C, Metcalf V (2011) Ocean
 501 acidification at high latitudes: potential effects on functioning of the antarctic bivalve
 502 *Laternula elliptica*. *Plos One* **6**: e16069

503 De Montaudouin X, Audemard C, Labourg P-J (1999) Does the slipper limpet (*Crepidula*
504 *fornicata*, L.) impair oyster growth and zoobenthos biodiversity? A revisited
505 hypothesis. *Journal of Experimental Marine Biology and Ecology* **235**: 105-124

506 Decottignies P, Beninger PG, Rincé Y, Riera P (2007) Trophic interactions between two
507 introduced suspension-feeders, *Crepidula fornicata* and *Crassostrea gigas*, are
508 influenced by seasonal effects and qualitative selection capacity. *Journal of*
509 *Experimental Marine Biology and Ecology* **342**: 231-241

510 Dickson AG, Millero FJ (1987) A comparison of the equilibrium constants for the
511 dissociation of carbonic acid in seawater media. *Deep Sea Research* **34**: 1733-1743

512 Dickson AG, Sabine CL, Christian JR (2007) Guide to best practices for ocean CO₂
513 measurements PICES special publication. North Pacific Marine Science
514 Organization, Sidney, British Columbia, pp 176

515 Diederich CM, Pechenik JA (2013) Thermal tolerance of *Crepidula fornicata* (Gastropoda)
516 life history stages from intertidal and subtidal subpopulations. *Marine Ecology-*
517 *Progress Series* **486**: 173-187

518 Dukes JS, Mooney HA (1999) Does global change increase the success of biological
519 invaders? *Trends in Ecology & Evolution* **14**: 135-139

520 Fabry VJ, Seibel BA, Feely RA, Orr JC (2008) Impacts of ocean acidification on marine
521 fauna and ecosystem processes. *Ices Journal of Marine Science* **65**: 414-432

522 Fernandez-Reiriz MJ, Range P, Alvarez-Salgado XA, Espinosa J, Labarta U (2012) Tolerance
523 of juvenile *Mytilus galloprovincialis* to experimental seawater acidification. *Marine*
524 *Ecology-Progress Series* **454**: 65-74

525 Fernandez-Reiriz MJ, Range P, Alvarez-Salgado XA, Labarta U (2011) Physiological
526 energetics of juvenile clams *Ruditapes decussatus* in a high CO₂ coastal ocean. *Marine*
527 *Ecology-Progress Series* **433**: 97-105

528 Fresard M, Boncoeur J (2006) Costs and benefits of stock enhancement and biological
529 invasion control: the case of the Bay of Brest scallop fishery. *Aquatic Living*
530 *Resources* **19**: 299-305

531 Gazeau F, Quiblier C, Jansen JM, Gattuso J-P, Middelburg JJ, Heip CHR (2007) Impact of
532 elevated CO₂ on shellfish calcification. *Geophysical Research Letters* **34**: 5

533 Guppy M, Withers P (1999) Metabolic depression in animals: physiological perspectives and
534 biochemical generalizations. *Biological Reviews of the Cambridge Philosophical*
535 *Society* **74**: 1-40

536 Gutowska MA, Melzner F, Langenbuch M, Bock C, Claireaux G, Pörtner H-O (2010) Acid-
537 base regulatory ability of the cephalopod (*Sepia officinalis*) in response to
538 environmental hypercapnia. *Journal of Comparative Physiology B* **180**: 323-335

539 Gutowska MA, Pörtner H-O, Melzner F (2008) Growth and calcification in the cephalopod
540 *Sepia officinalis* under elevated seawater pCO₂. *Marine Ecology-Progress Series* **373**:
541 303-309

542 Hale R, Calosi P, McNeill L, Mieszkowska N, Widdicombe S (2011) Predicted levels of
543 future ocean acidification and temperature rise could alter community structure and
544 biodiversity in marine benthic communities. *Oikos* **120**: 661-674

545 Hellmann JJ, Byers JE, Bierwagen BG, Dukes JS (2008) Five potential consequences of
546 climate change for invasive species. Cinco Consecuencias Potenciales del Cambio
547 Climático para Especies Invasoras. *Conservation Biology* **22**: 534-543

548 Hiebenthal C, Philipp EE, Eisenhauer A, Wahl M (2013) Effects of seawater pCO₂ and
549 temperature on shell growth, shell stability, condition and cellular stress of Western
550 Baltic Sea *Mytilus edulis* (L.) and *Arctica islandica* (L.). *Marine Biology* **160**: 2073-
551 2087

552 Hofmann GE, Todgham AE (2010) Living in the now: physiological mechanisms to tolerate a
 553 rapidly changing environment. *Annual Review of Physiology* **72**: 127-145
 554 Kroeker KJ, Gambi MC, Micheli F (2013a) Community dynamics and ecosystem
 555 simplification in a high-CO₂ ocean. *Proceedings of the National Academy of Sciences*
 556 **110**: 12721-12726
 557 Kroeker KJ, Kordas RL, Crim R, Hendriks IE, Ramajo L, Singh GS, Duarte CM, Gattuso J-P
 558 (2013b) Impacts of ocean acidification on marine organisms: quantifying sensitivities
 559 and interaction with warming. *Global Change Biology* **19**: 1884-1896
 560 Langenbuch M, Bock C, Leibfritz D, Pörtner H-O (2006) Effects of environmental
 561 hypercapnia on animal physiology: A C-13 NMR study of protein synthesis rates in
 562 the marine invertebrate *Sipunculus nudus*. *Comparative Biochemistry and Physiology*
 563 *A-Molecular & Integrative Physiology* **144**: 479-484
 564 Lannig G, Eilers S, Pörtner H-O, Sokolova IM, Bock C (2010) Impact of ocean acidification
 565 on energy metabolism of oyster, *Crassostrea gigas* - Changes in metabolic pathways
 566 and thermal response. *Marine Drugs* **8**: 2318-2339
 567 Lenz M, da Gama BAP, Gerner NV, Gobin J, Groner F, Harry A, Jenkins SR, Kraufvelin P,
 568 Mummelthei C, Sareyka J, Xavier EA, Wahl M (2011) Non-native marine
 569 invertebrates are more tolerant towards environmental stress than taxonomically
 570 related native species: Results from a globally replicated study. *Environmental*
 571 *Research* **111**: 943-952
 572 Lewis E, Wallace DWR (1998) Program developed for CO₂ system calculations. Carbon
 573 Dioxide Information Analysis Center, Oak Ridge National Laboratory, U.S.
 574 Department of Energy

575 Liu W, He M (2012) Effects of ocean acidification on the metabolic rates of three species of
 576 bivalve from southern coast of China. *Chinese Journal of Oceanology and Limnology*
 577 **30**: 206-211

578 Mackenzie CL, Ormondroyd GA, Curling SF, Ball RJ, Whiteley NM, Malham SK (2014)
 579 Ocean warming, more than acidification, reduces shell strength in a commercial
 580 shellfish species during food limitation. *Plos One* **9**: e86764

581 Marchant HK, Calosi P, Spicer JJ (2010) Short-term exposure to hypercapnia does not
 582 compromise feeding, acid-base balance or respiration of *Patella vulgata* but
 583 surprisingly is accompanied by radula damage. *Journal of the Marine Biological*
 584 *Association of the United Kingdom* **90**: 1379-1384

585 Marshall DJ, Perissinotto R, Holley JF (2003) Respiratory responses of the mysid
 586 *Gastrosaccus brevifissura* (Peracarida : Mysidacea), in relation to body size,
 587 temperature and salinity. *Comparative Biochemistry and Physiology A-Molecular &*
 588 *Integrative Physiology* **134**: 257-266

589 Martin S, Thouzeau G, Chauvaud L, Jean F, Guérin L (2006) Respiration, calcification, and
 590 excretion of the invasive slipper limpet, *Crepidula fornicata* L.: Implications for
 591 carbon, carbonate, and nitrogen fluxes in affected areas. *Limnology & Oceanography*
 592 **51**: 1996-2007

593 Martin S, Thouzeau G, Richard M, Chauvaud L, Jean F, Clavier J (2007) Benthic community
 594 respiration in areas impacted by the invasive mollusk *Crepidula fornicata*. *Marine*
 595 *Ecology-Progress Series* **347**: 51-60

596 Matozzo V, Chinellato A, Munari M, Bressan M, Marin MG (2013) Can the combination of
 597 decreased pH and increased temperature values induce oxidative stress in the clam
 598 *Chamelea gallina* and the mussel *Mytilus galloprovincialis*? *Marine Pollution Bulletin*
 599 **72**: 34-40

600 Matozzo V, Chinellato A, Munari M, Finos L, Bressan M, Marin MG (2012) First evidence of
 601 immunomodulation in bivalves under seawater acidification and increased
 602 temperature. *Plos One* **7**: e33820
 603 Mehrbach C, Culberso.Ch, Hawley JE, Pytkowic RM (1973) Measurement of apparent
 604 dissociation-constants of carbonic-acid in seawater at atmospheric-pressure.
 605 *Limnology & Oceanography* **18**: 897-907
 606 Melatunan S, Calosi P, Rundle SD, Moody AJ, Widdicombe S (2011) Exposure to elevated
 607 temperature and $p\text{CO}_2$ reduces respiration rate and energy status in the periwinkle
 608 *Littorina littorea*. *Physiological and Biochemical Zoology* **84**: 583-594
 609 Melzner F, Gutowska MA, Langenbuch M, Dupont S, Lucassen M, Thorndyke MC, Bleich
 610 M, Pörtner H-O (2009) Physiological basis for high CO_2 tolerance in marine
 611 ectothermic animals: pre-adaptation through lifestyle and ontogeny? *Biogeosciences* **6**:
 612 2313-2331
 613 Melzner F, Stange P, Trubenbach K, Thomsen J, Casties I, Panknin U, Gorb SN, Gutowska
 614 MA (2011) Food supply and seawater $p\text{CO}_2$ impact calcification and internal shell
 615 dissolution in the blue mussel *Mytilus edulis*. *Plos One* **6**: e24223
 616 Metzger R, Sartoris FJ, Langenbuch M, Pörtner H-O (2007) Influence of elevated CO_2
 617 concentrations on thermal tolerance of the edible crab *Cancer pagurus*. *Journal of*
 618 *Thermal Biology* **32**: 144-151
 619 Michaelidis B, Ouzounis C, Paleras A, Pörtner H-O (2005) Effects of long-term moderate
 620 hypercapnia on acid-base balance and growth rate in marine mussels *Mytilus*
 621 *galloprovincialis*. *Marine Ecology-Progress Series* **293**: 109-118
 622 Morley SA, Peck LS, Miller AJ, Pörtner HO (2007) Hypoxia tolerance associated with
 623 activity reduction is a key adaptation for *Laternula elliptica* seasonal energetics.
 624 *Oecologia* **153**: 29-36

625 Navarro JM, Torres R, Acuña K, Duarte C, Manriquez PH, Lardies M, Lagos NA, Vargas C,
 626 Aguilera V (2013) Impact of medium-term exposure to elevated $p\text{CO}_2$ levels on the
 627 physiological energetics of the mussel *Mytilus chilensis*. *Chemosphere* **90**: 1242-1248
 628 Newell RC, Kofoed LH (1977) Adjustment of components of energy-balance in gastropod
 629 *Crepidula fornicata* in response to thermal acclimation. *Marine Biology* **44** : 275-286
 630 Noisette F, Duong G, Six C, Davoult D, Martin S (2013) Effects of elevated $p\text{CO}_2$ on the
 631 metabolism of a temperate rhodolith *Lithothamnion corallioides* grown under different
 632 temperatures. *Journal of Phycology* **49**: 746-757
 633 Norkko J, Thrush SF (2006) Ecophysiology in environmental impact assessment: implications
 634 of spatial differences in seasonal variability of bivalve condition. *Marine Ecology*
 635 *Progress Series* **326**: 175-186
 636 Occhipinti-Ambrogi A (2007) Global change and marine communities: Alien species and
 637 climate change. *Marine Pollution Bulletin* **55**: 342-352
 638 Pansch C, Schaub I, Havenhand J, Wahl M (2014) Habitat traits and food availability determine the response of marine
 639 invertebrates to ocean acidification. *Global Change Biology* **20**: 265-277
 640 Pansch C, Schaub I, Havenhand J, Wahl M (2014) Habitat traits and food availability
 641 determine the response of marine invertebrates to ocean acidification. *Global Change*
 642 *Biology* **20**: 265-277
 643 Parsons TR, Takahashi M, Hargrave B (1984) Biological oceanographic processes (3rd ed). In:
 644 (Eds) BH (ed), Oxford, pp 330
 645 Peck LS, Clark MS, Morley SA, Massey A, Rossetti H (2009) Animal temperature limits and
 646 ecological relevance: effects of size, activity and rates of change. *Functional Ecology*
 647 **23**: 248-256

648 Peck LS, Convey P, Barnes DKA (2006) Environmental constraints on life histories in
649 antarctic ecosystems: Tempos, timings and predictability, *Biological Reviews* **81**: 75-
650 109

651 Peck LS, Morley SA, Richard J, Clark MS (2014) Acclimation and thermal tolerance in
652 Antarctic marine ectotherms. *Journal of Experimental Biology* **217**: 16-22

653 Peck LS, Pörtner H-O, Hardewig I (2002) Metabolic demand, oxygen supply, and critical
654 temperatures in the Antarctic bivalve *Laternula elliptica*. *Physiological and*
655 *Biochemical Zoology* **75**: 123-133

656 Peck LS, Souster T, Clark MS (2013) Juveniles are more resistant to warming than adults in 4
657 species of Antarctic marine invertebrates. *PLoS One* **8**: e66033

658 Peck LS, Webb KE, Bailey DM (2004) Extreme sensitivity of biological function to
659 temperature in Antarctic marine species. *Functional Ecology* **18**: 625-630

660 Pörtner H-O (2008) Ecosystem effects of ocean acidification in times of ocean warming: a
661 physiologist's view. *Marine Ecology-Progress Series* **373**: 203-217

662 Pörtner H-O (2012) Integrating climate-related stressor effects on marine organisms: unifying
663 principles linking molecule to ecosystem-level changes. *Marine Ecology-Progress*
664 *Series* **470**: 273-290

665 Pörtner H-O, Langenbuch M, Michaelidis B (2005) Synergistic effects of temperature
666 extremes, hypoxia, and increases in CO₂ on marine animals: from Earth history to
667 global change. *Journal of Geophysical Research-Oceans* **110**: C09S10

668 Pörtner H-O, Peck LS, Hirse T (2006) Hyperoxia alleviates thermal stress in the Antarctic
669 bivalve, *Laternula elliptica*: evidence for oxygen limited thermal tolerance. *Polar*
670 *Biology* **29**: 688-693

671 R Core Team (2013) R: a language and environment for statistical computing. R Foundation
672 for Statistical Computing, Vienna, Austria

673 Range P, Chicharo MA, Ben-Hamadou R, Pilo D, Fernandez-Reiriz MJ, Labarta U, Marin
674 MG, Bressan M, Matozzo V, Chinellato A, Munari M, El Menif NT, Dellali M,
675 Chicharo L (2014) Impacts of CO₂-induced seawater acidification on coastal
676 Mediterranean bivalves and interactions with other climatic stressors. *Regional*
677 *Environmental Change* **14** (Suppl 1): S19-S30

678 Richard J, Morley SA, Deloffre J, Peck LS (2012) Thermal acclimation capacity for four
679 Arctic marine benthic species. *Journal of Experimental Marine Biology and Ecology*
680 **424**: 38-43

681 Sanders MB, Bean TP, Hutchinson TH, Le Quesne WJF (2013) Juvenile king scallop, *Pecten*
682 *maximus*, is potentially tolerant to low levels of ocean acidification when food is
683 unrestricted. *Plos One* **8**: e74118

684 Schalkhausser B, Bock C, Stemmer K, Brey T, Pörtner H-O, Lannig G (2013) Impact of
685 ocean acidification on escape performance of the king scallop, *Pecten maximus*, from
686 Norway. *Marine Biology* **160**: 1995-2006

687 Shirayama Y, Thornton H (2005) Effect of increased atmospheric CO₂ on shallow water
688 marine benthos. *Journal of Geophysical Research: Oceans* **110** (C09S08): 1-8

689 Sokolova IM, Pörtner H-O (2003) Metabolic plasticity and critical temperatures for aerobic
690 scope in a eurythermal marine invertebrate (*Littorina saxatilis*, Gastropoda :
691 Littorinidae) from different latitudes. *Journal of Experimental Biology* **206**: 195-207

692 Solomon S, Quin D, Manning M, Chen Z, Marquis M, Averyt K, Tignor M, Miler H (2007)
693 Contribution of working group I to the fourth assessment report of the
694 Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge,
695 pp 996

696 Thomsen J, Casties I, Pansch C, Kortzinger A, Melzner F (2013) Food availability outweighs
 697 ocean acidification effects in juvenile *Mytilus edulis*: laboratory and field experiments.
 698 *Global Change Biology* **19**: 1017-1027

699 Thomsen J, Gutowska MA, Saphorster J, Heinemann A, Trubenbach K, Fietzke J, Hiebenthal
 700 C, Eisenhauer A, Kortzinger A, Wahl M, Melzner F (2010) Calcifying invertebrates
 701 succeed in a naturally CO₂-rich coastal habitat but are threatened by high levels of
 702 future acidification. *Biogeosciences* **7**: 3879-3891

703 Thomsen J, Melzner F (2010) Moderate seawater acidification does not elicit long-term
 704 metabolic depression in the blue mussel *Mytilus edulis*. *Marine Biology* **157**: 2667-
 705 2676

706 Thouzeau G, Chauvaud L, Grall J, Guérin L (2000) Rôle des interactions biotiques sur le
 707 devenir du pré-recrutement et la croissance de *Pecten maximus* (L.) en rade de Brest.
 708 *Comptes Rendus de l'Académie des Sciences - Series III - Sciences de la Vie* **323**: 815-
 709 825

710 Tomanek L, Zuzow MJ, Ivanina AV, Beniash E, Sokolova IM (2011) Proteomic response to
 711 elevated pCO₂ level in eastern oysters, *Crassostrea virginica*: evidence for oxidative
 712 stress. *Journal of Experimental Biology* **214**: 1836-1844

713 von Bertalanffy L (1951) Metabolic types and growth types. *The American Naturalist* **85**:
 714 111-117

715 Watson S-A, Peck LS, Tyler PA, Southgate PC, Tan KS, Day RW, Morley SA (2012) Marine
 716 invertebrate skeleton size varies with latitude, temperature and carbonate saturation:
 717 implications for global change and ocean acidification. *Global Change Biology* **18**:
 718 3026-3038

719 Willson LL, Burnett LE (2000) Whole animal and gill tissue oxygen uptake in the Eastern
 720 oyster, *Crassostrea virginica*: effects of hypoxia, hypercapnia, air exposure, and

infection with the protozoan parasite *Perkinsus marinus*. *Journal of Experimental Marine Biology and Ecology* **246**: 223-240

Wood HL, Spicer JJ, Lowe DM, Widdicombe S (2010) Interaction of ocean acidification and temperature; the high cost of survival in the brittlestar *Ophiura ophiura*. *Marine Biology* **157**: 2001-2013

Wood HL, Spicer JJ, Widdicombe S (2008) Ocean acidification may increase calcification rates, but at a cost. *Proceedings of the Royal Society B-Biological Sciences* **275**: 1767-1773

FIGURES CAPTIONS

Figure 1: Cumulated mortalities during the temperature increase. Males are represented on the graph on the top and females are on the graph in the bottom. The greyscale represent the different $p\text{CO}_2$ levels in which *C. fornicata* individuals where held during the experiment.

Figure 2: Mean (\pm SE) conditions indices at the beginning (black bar), and at the end of the experiment for *C. fornicata* females (white bars) and males (grey bars) in the different $p\text{CO}_2$.
 $27 > N > 10$

Figure 3: Respiration rates as a function of increasing temperature in each $p\text{CO}_2$ treatment, for *C. fornicata* males (top, triangles) and females (bottom, circles). Detailed statistical analyses relative to the regressions can be found in Table 3.

Figure 4: Mean (\pm SE) Q_{10} values for *C. fornicata* females (white bars) and males (grey bars) in the different $p\text{CO}_2$ treatments. $N = 3$

TABLES

Table 1: Mean (\pm standard error, SE) carbonate chemistry parameters for each $p\text{CO}_2$ treatment. pH (on the total scale, pH_T) was measured daily and total alkalinity (A_T) was measured every 3 weeks. Other parameters were calculated with CO2sys software. $p\text{CO}_2$: CO_2 partial pressure; Ω_{Ar} : saturation state of seawater with respect to aragonite.

$p\text{CO}_2$ treatment	pH_T	$p\text{CO}_2$ (μatm)	Ω_{Ar}	A_T ($\mu\text{Eq kg}^{-1}$ SW)
	n = 69	n = 69	n = 69	n = 76
380 μatm	8.13 ± 0.01	324 ± 8	2.72 ± 0.06	2333 ± 1
550 μatm	7.89 ± 0.01	619 ± 16	1.69 ± 0.04	2334 ± 2
750 μatm	7.75 ± 0.01	873 ± 20	1.28 ± 0.03	2335 ± 2
1000 μatm	7.66 ± 0.01	1138 ± 65	1.05 ± 0.02	2334 ± 2

Table 2: Summary of two-way ANOVAs testing the effects of $p\text{CO}_2$, sex and the interaction of these two factors on the final condition indices (CI) and the Q_{10} values determined for *C. fornicata* males and females in the different $p\text{CO}_2$ conditions (380, 550, 750 and 1000 μatm). Bold numbers indicate significant level greater than 95%.

	df	CI		Q_{10}	
		F-value	p-value	F-value	p-value
$p\text{CO}_2$	3	1.245	0.295	0.657	0.590
sex	1	2.472	0.118	6.124	0.025
$p\text{CO}_2 \times \text{sex}$	3	1.371	0.254	2.293	0.117

Table 3: Relationships between *C. fornicata* male and female respiration rates and temperature in each $p\text{CO}_2$ treatment

	$p\text{CO}_2$	Regression equation	n	R	R^2	F	p
males	380	$y = 3.691 x - 34.455$	42	0.60	0.37	22.97	< 0.001
	550	$y = 2.993 x - 18.461$	42	0.46	0.21	10.56	0.002
	750	$y = 2.406 x - 4.543$	41	0.40	0.16	7.55	0.009
	1000	$y = 3.701 x - 41.556$	41	0.56	0.31	17.37	< 0.001
females	380	$y = 1.826 x - 7.635$	42	0.49	0.24	12.72	< 0.001
	550	$y = 1.585 x - 4.218$	42	0.55	0.30	16.89	< 0.001
	750	$y = 2.637 x - 26.240$	42	0.63	0.40	26.66	< 0.001
	1000	$y = 1.442 x + 3.435$	42	0.37	0.14	6.26	0.017